

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending April 11, 2015

Prepared by the Office of Infectious Disease Epidemiology and Outbreak Response Infectious Disease Bureau – Prevention and Health Promotion Administration Maryland Department of Health and Mental Hygiene

The data presented in this document are provisional and subject to change as additional reports come in.

SUMMARY

During the week ending April 11, 2015, influenza-like illness (ILI) intensity in Maryland was MINIMAL and there was REGIONAL geographic activity. The proportion of outpatient visits for ILI reported by Sentinel Providers climbed, while the proportion reported by Maryland Emergency Departments was unchanged. The proportion of MRITS respondents reporting ILI decreased slightly. The percentage of specimens testing positive at clinical laboratories was down, but remained elevated, with a majority of positives being influenza type B. The number of specimens testing positive at the DHMH lab remained the same. A total of 36 influenza-associated hospitalizations were reported. There was 1 respiratory outbreak. Nationally, influenza activity continued to decline.

ILI Intensity Levels

Minimal

Low

Moderate


High

Influenza Geographic Spread
No Activity
Sporadic
Local
✓ Regional
Widespread

Click here to visit our influenza surveillance web page

ILINet Sentinel Providers


Twenty-eight sentinel providers reported a total of 10,245 visits this week. Of those, 159 (1.6%) were visits for ILI. This is below the Maryland baseline of 2.0%.



ILI Visits To Sentinel Providers By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	33 (21%)	34 (24%)	1336 (24%)
Age 5-24	47 (30%)	50 (36%)	2200 (40%)
Age 25-49	48 (30%)	34 (24%)	1190 (22%)
Age 50-64	21 (13%)	13 (9%)	517 (9%)
Age ≥ 65	10 (6%)	8 (6%)	253 (5%)
Total ILI Visits	159 (100%)	139 (100%)	5496 (100%)

Visits to Emergency Departments for ILI

Emergency Departments in Maryland reported a total of 49,126 visits this week through the ESSENCE surveillance system. Of those, 991 (2.0%) were visits for ILI.



ILI Visits To Emergency Departments By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	260 (26%)	269 (28%)	7982 (25%)
Age 5-24	279 (28%)	299 (31%)	9918 (31%)
Age 25-49	259 (26%)	264 (27%)	8578 (26%)
Age 50-64	122 (12%)	74 (8%)	3435 (11%)
Age ≥ 65	71 (7%)	61 (6%)	2529 (8%)
Unknown			3 (<1%)
Total ILI Visits	991 (100%)	967 (100%)	32445(100%)

Other states' influenza information:

Delaware http://dhss.delaware.gov/dph/epi/influenzahome.html

District of Columbia http://doh.dc.gov/service/influenza

Pennsylvania http://www.portal.state.pa.us/portal/server.pt/community/influenza_(flu)/14161

Virginia http://www.vdh.state.va.us/Epidemiology/flu/


West Virginia http://dhhr.wv.gov/oeps/disease/flu/Pages/fluSurveillance.aspx


Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending April 11, 2015

Community-based Influenza Surveillance (MRITS and Google Flu Trends)


MRITS is the Maryland Resident Influenza Tracking System, a weekly survey for influenza-like illness (ILI). A total of 620 residents responded to the MRITS survey this week. Of those, 6 (1.0%) reported having ILI and missing a cumulative 6 days of regular daily activities. According to Google Flu Trends, influenza activity is "Low" in Maryland.






Clinical Laboratory Influenza Testing

Twenty-seven clinical laboratories reported performing 1,164 influenza diagnostic tests, mostly rapid influenza diagnostic tests (RIDTs). Of those, 163 (14.0%) were positive for influenza. Of those testing positive, 9 (5.5%) were influenza Type A and 154 (94.5%) were influenza Type B. The reliability of RIDTs depends largely on the conditions under which they are used. False-positive (and true-negative) results are more likely to occur when the disease prevalence in the community is low, which is generally at the beginning and end of the influenza season and during the summer.




Positive Rapid Flu Tests by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A	9 (6%)	58 (30%)	13887 (85%)
Туре В	154 (94%)	133 (70%)	2377 (15%)
Total	163 (100%)	191 (100%)	16264(100%)



State Laboratories Administration Influenza Testing

The DHMH Laboratories Administration performed a total of 76 PCR tests for influenza and 25 (32.9%) specimens tested positive. Of those, 1 (4.0%) was Type A (H1N1), 2 (8.0%) were Type A (H3N2), and 22 (88.0%) were Type B. PCR testing is more reliable than RIDT. The DHMH testing identifies subtypes of influenza A, information that is not available from the RIDT results. The table below summarizes results by type and subtype.



Positive PCR Tests by Type (Subtype)	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A (H1)	1 (4%)		3 (<1%)
Type A (H3)	2 (8%)	1 (4%)	2383 (95%)
Туре В	22 (88%)	24 (96%)	125 (5%)
A (H3) – B Dual Infection			2 (<1%)
Total	25 (100%)	25 (100%)	2513 (100%)

Where to get an influenza vaccination


Interested in getting a flu vaccine for the 2014-15 influenza season? Go to http://dhmh.maryland.gov/flumd/SitePages/getvaccinated.aspx and click on your county/city of residence. You will be redirected to your local health department website for local information on where to get your flu vaccine.

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending April 11, 2015

Influenza-associated Hospitalizations


For this week, hospitals in Maryland reported 36 influenza-associated hospitalizations. (A person with an overnight hospital stay along with a positive influenza test of any kind, e.g. RIDT or PCR, is considered an "influenza-associated hospitalization" for purposes of influenza surveillance.)



Influenza- Associated Hospitalizations by Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	3 (8%)	4 (8%)	174 (5%)
Age 5-17		3 (6%)	89 (3%)
Age 18-24		1 (2%)	79 (2%)
Age 25-49	7 (19%)	8 (16%)	380 (11%)
Age 50-64	12 (33%)	12 (24%)	588 (17%)
Age ≥ 65	14 (39%)	23 (45%)	2214 (63%)
Total	36 (100%)	51 (100%)	3524 (100%)

Outbreaks of Respiratory Disease

There was 1 respiratory outbreak reported to DHMH this week. (Disease outbreaks of any kind are reportable in Maryland. Respiratory outbreaks may be reclassified once a causative agent is detected, e.g. from ILI to influenza.)



Respiratory Outbreaks by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Influenza	-	3 (75%)	126 (70%)
Influenza-like Illness			36 (20%)
Pneumonia	1 (100%)	1 (25%)	18 (10%)
Other Respiratory			
Total	1 (100%)	4 (100%)	180 (100%)

National Influenza Surveillance (CDC)

During week 14 (April 5-11, 2015), influenza activity continued to decrease in the United States.

- Viral Surveillance: Of 11,189 specimens tested and reported by U.S. World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NREVSS) collaborating laboratories during week 14, 1,076 (9.6%) were positive for influenza.
- O Pneumonia and Influenza Mortality: The proportion of deaths attributed to pneumonia and influenza (P&I) was below the epidemic threshold.
- Influenza-associated Pediatric Deaths: No influenza-associated pediatric deaths were reported.
- Influenza-associated Hospitalizations: A cumulative rate for the season of 62.4 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported.
- Outpatient Illness Surveillance: The proportion of outpatient visits for influenza-like illness (ILI) was 1.8%, which is below the national baseline of 2.0%. Three regions reported ILI at or above region-specific baseline levels. Puerto Rico and six states experienced low ILI activity; New York City and 44 states experienced minimal ILI activity; and the District of Columbia and the U.S. Virgin Islands had insufficient data.
- Geographic Spread of Influenza: The geographic spread of influenza in three states was reported as widespread; Guam and 14 states reported regional activity; the District of Columbia and 19 states reported local activity; and Puerto Rico, the U.S. Virgin Islands, and 14 states reported sporadic activity.

